

## **Guidelines related to your Facial Surgery**

Dear patient,

questions related to facial surgery procedures often arise before your surgery. To help you to prepare for your operation and the time afterwards, we have put some guidelines together for you.

### **Preparation for your operation**

- If you undergo your operation under general anaesthesia you will have to be fasting. Please discuss the start and duration of the fasting period with the anaesthesiologist during your anaesthetic consultation.
- If you wish to buy additional private insurance (“Folgekostenversicherung”), you will have to do that at least one day before your surgery. Make sure you choose the right type of insurance!
- Stop taking blood thinners, e.g. Aspirin®, ASS or Cumarine at least 2 weeks before your operation. If you need to take the medication on a regularly basis please discuss with your family doctor/internist before your operation how/when to stop or "bridge" your medication.
- Nicotine should be avoided at least 2 weeks before and after your operation, preferably 6 weeks before and after your operation. This also applies to chewing tobacco, water pipes, nicotine patches and nicotine gum.
- You should avoid alcohol and excessive consumption of coffee and chocolate for at least 1 week before and after your operation.
- Please refrain from wearing face creams, make-up, eyelash extensions and contact lenses on the day of your operation.
- No metal may be worn on the body during the operation, therefore remove jewellery, earrings, piercings etc. beforehand.

### **After your operation**

- Get someone to pick you up after your operation. Under no circumstances you should drive yourself! Furthermore someone should be at home with you at least for the first night.
- Drink plenty and regularly in small amounts after your operation.

### **Aftercare**

- After your operation you should not just lie in bed or on the couch. The best prophylaxis of thrombosis is exercise! Therefore, go for regular walks, ideally accompanied.
- Bruising and/or swelling after your operation is normal and will subside within a few weeks. You can cool your face. However, please avoid direct contact of ice with the skin or eyes. If autologous fat has been injected, the injected areas must not be cooled!
- You will have to sleep on your back for the first 2 weeks after your operation. Sleep with the headboard raised for the first few nights or use several pillows if your bed is not adjustable.

- After eyelid surgery, the eyes can react sensitively to wind and sunlight. Wearing sunglasses can be helpful in these cases. And buy yourself new make-up. Old facial powder, eyeliner pencils, mascara etc. are often contaminated with bacteria. These bacteria should not get into fresh wounds.
- Avoid working with your head bent forwards for the first few days after the operation. This applies in particular to rhinoplasties. Avoid blood pressure spikes and high pulse rates, e.g. due to overexertion, sport or sexual intercourse.
- Leave the tapes on the wound until the follow-up appointment. The silicone splints and nasal splint must never be removed without authorisation by Dr. Sattler after rhinoplasties! After your operation on the nose, you must not blow your nose or pull it up. Only dab the nose carefully with a handkerchief.
- The stitches are usually removed about 1 week after the operation, and in the case of operations on the nose, the silicone splints in the nose and the outer aluminium splint are also removed.
- You can start scar care approximately 2 weeks after your operation. We recommend a silicone-based scar gel. Fresh scars do not tolerate UV light. Direct sunlight or visits to sun beds are therefore not recommended for at least a few months. Some scar gels contain sun protection factors are very useful.
- Sauna, steam baths, swimming pools and whirlpools should be avoided for at least 6 weeks. The same applies to bath tubs, bathing in lakes, the sea or public swimming pools.
- Eat healthy and make sure you have an adequate intake of Proteins, Vitamin C and Zinc. Do not stay fasting for health or religious reasons after your operation, since this could impair wound healing. Eating pineapple helps against swelling. The pineapple enzyme (Bromelain) can also be taken in tablet form. Heparin ointment helps against bruising and diclofenac gel against swelling, pain and muscle aches. Of course, these substances should not be used if you suffer from any allergies related to these substances!
- Dr. Sattler cannot issue a sick note ("AU") for work after your operation. Please contact your family doctor in this regard. However, the German law says that you have to take holiday leave if you are unable to work after an aesthetic procedure.

Give yourself and your body a break and be patient. Make sure to attend the check-up appointments with Dr. Sattler and follow the guidelines stated above. In urgent emergencies outside our consultation hours, please contact a plastic surgery emergency clinic, e.g. at the UKE in Hamburg.

*Your Plastic Surgery Team Dr. Sattler*

**I have taken note of the guidelines and will comply with them.**

---

**Place, Date, Signature of the Patient**